
Time: 13:00-18:00

Instructions:

- During the exam you MAY NOT use textbooks, class notes, or any other supporting material.
- Use of calculators is permitted for performing calculations. The use of graphic or programmable features is NOT permitted.
- In all of your solutions, give explanations to clearly show your reasoning. Points may be deducted for unclear solutions even if the answer is correct.
- Use natural language when appropriate, not just mathematical symbols.
- Write clearly and legibly.
- Where applicable, indicate your final answer clearly by putting A BOX around it.

Grades: There are 6 problems. A maximum of 5 points can be awarded for each problem solved. At least 15 points are necessary for the grade E. The problems are not ordered according to difficulty.

1. Let A be the matrix

$$A = \begin{pmatrix} 2 & 5 & -4 \\ -2 & k-8 & 6 \\ 4 & 9-k & k-7 \end{pmatrix}.$$

depending on the parameter $k \in \mathbb{R}$.

- (a) Compute the determinant $|A|$ as a function of k . (2p)
- (b) Determine the values of k for which the matrix A is not invertible. (1p)
- (c) Solve the system of linear equations

$$\begin{cases} 2x + 5y - 4z = 1 \\ -2x - 8y + 6z = 0 \\ 4x + 9y - 7z = 2 \end{cases}$$

in the variables x , y and z using Gaussian elimination. (2p)

2. Consider the function $f(x) = \sqrt[3]{x}$.

- (a) Compute the Taylor polynomial $T_2(x)$ of order 2 of $f(x)$ about $x = 8$. (3p)
- (b) Use the answer from part (a) to estimate $\sqrt[3]{6}$. (2p)

3. (a) Compute the primitive $\int \frac{\ln(t)}{5t} \sqrt{1 + (\ln(t))^2} dt$ (as a function of t). (2p)

(b) Find a number a for which $\int_1^a \frac{1}{\sqrt{x}} + 1 dx = 5$ (3p).

4. Consider the function $f(x, y) = e^{xy-y}$ defined on the closed, bounded set

$$D = \{(x, y) \in \mathbb{R}^2 \mid x^2 - 2x - 1 \leq y \leq x - 1\}.$$

- (a) Determine the coordinates of the points of intersection of $\{y = x - 1\}$ with $\{y = x^2 - 2x - 1\}$. (1p)
- (b) Determine the critical points of f and compute the value of f at those points. (2p)
- (c) Determine the maximal and the minimal values of f on D . (2p)

5. (a) Compute

$$\lim_{x \rightarrow +\infty} \left(\frac{x^2}{x+2} - \frac{x^2+1}{x-3} \right) \quad (3p)$$

(b) Find a number a for which the following function is continuous for all x :

$$f(x) = \begin{cases} ax & x \geq 2 \\ x^2 + a & x < 2 \end{cases} \quad (2p)$$

6. Consider the function $f(x) = \frac{x}{x^2 - 3x + 2}$.

- (a) Determine the domain of definition of f . (1p)
- (b) Determine the local extreme points of f . (3p)
- (c) Determine where f is increasing and where f is decreasing. (1p)

Formulas

The Taylor polynomial $T_n(x)$ of order n of the function $f(x)$ at $x = x_0$ is

$$T_n(x) = f(x_0) + \frac{f'(x_0)}{1!}(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n.$$

The solutions of the equation $ax^2 + bx + c = 0$ are given by $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ if $b^2 - 4ac \geq 0$.

GOOD LUCK!