

---

Please **READ CAREFULLY** the general instructions:

- During the exam you CAN NOT use any textbook, class notes, or any other supporting material.
  - Calculators are **not allowed** during the exam.
  - In all your solutions show your reasoning, explaining carefully what you are doing. Justify your answers.
  - Use natural language, not just mathematical symbols.
  - Use clear and legible writing. Write preferably with a ball-pen or a pen (black or dark blue ink).
  - **Do not write two exercises on the same page.**
  - The written exam gives up to 24 points. A minimum of 15 points (including bonus points) guarantees a pass grade. A minimum of 21 points in the written exam guarantees access to the oral exam.
  - The list of the anonymous codes of those who qualify for the oral exam will be published on the course page by Friday 19/1 the latest. The oral exam will take place in Albano premises on **Thursday 25/1** (and eventually also Friday 26/1).
- 

1. Determine if the following sets are countable or not:

- (a) [2pt] The family of all functions from  $\mathbb{N}$  to  $\{0, 1, 2\}$  i.e.  $\{f : f : \mathbb{N} \rightarrow \{0, 1, 2\}\}$ .
- (b) [1pt] The set of all affine lines in the plane, which are perpendicular to the vector  $(-2, 1)$  and have a nonempty intersection with the set  $\{(x, y) : x = 0, y \in \mathbb{Q}\}$ .

2. Let  $(X, d)$  be a metric space, and let  $E, K$  be two subsets of  $X$ .

- (a) [1pt] Define what it means for a set  $K \subset X$  to be compact.
- (b) [1pt] Define what it means to be a limit point of  $E$ .
- (c) [2pt] Let  $E \subset K$  and assume that  $K$  is compact. Prove in detail that  $E$  has no limit point in  $K$  if, and only if  $E$  is finite.

3. [3pt] Let  $\{x_n\}_{n=1}^\infty$  be a sequence in a complete metric space  $(X, d)$ , for which there exists a positive constant  $C < 1$  such that for all  $n \geq 2$ ,

$$d(x_{n+1}, x_n) \leq C d(x_n, x_{n-1}).$$

Prove that the sequence  $\{x_n\}_{n=1}^\infty$  is convergent.

4. (a) [3pt] Given  $n \in \mathbb{N}$ , let  $f_n(x) = x(1 - x^2)^n$  defined for  $x \in [0, 1]$ . Study the pointwise and uniform convergence of the sequence  $\{f_n\}_{n \geq 1}$  on  $[0, 1]$ .
- (b) [2pt] Let  $\{f_n\}_{n \geq 1}$  be a sequence of real-valued bounded functions defined on  $[0, 1]$  for which

$$\limsup_n (\|f_n\|_\infty (n(\log n)^2)) = C \in [0, \infty),$$

where  $\|f_n\|_\infty = \sup_{x \in [0, 1]} |f_n(x)|$ . Prove that the series  $\sum f_n$  converges uniformly.

5. [3pt] Consider the following system of equations

$$\begin{cases} xu + y^2v = 0 \\ xv^4 - y^3u^5 = 0. \end{cases}$$

Show that near the point  $(x_0, y_0, u_0, v_0) = (1, -1, 0, 1)$  we can write  $u, v$  as a  $\mathcal{C}^1$  function of  $x, y$ . Calculate  $\frac{\partial v}{\partial y}(1, -1)$ .

Please turn page  $\longrightarrow$

6. Determine which of the following statements are true and which are false. Explain your reasoning, by giving a proof or a counterexample to each statement. Each answer is graded over one point. Please summarise your answers in the form that you can find in the cover sheet.
- i. There exists two increasing functions  $g, h: [0, 2024] \rightarrow \mathbb{R}$  such that the function  $f(x) = g(x) - h(x)$  has an uncountable number of discontinuities.
  - ii. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is given by  $f(x) = 1/(2+x^2)$  then, for every closed set  $K \subset \mathbb{R}$ ,  $f^{-1}(K)$  is a closed set of  $\mathbb{R}$ .
  - iii. Let  $a < b$  be two real numbers. There exist two bounded functions  $f, g: [a, b] \rightarrow \mathbb{R}$  for which  $\int_a^b (f+g)dx > \int_a^b fdx + \int_a^b gdx$ .
  - iv. If the power series with complex coefficients  $\sum_{k \geq 0} a_k z^k$  converges for  $z = 5i$ , it also converges for  $z = 1 - 2i$ .
  - v. For any real-valued continuous function  $f: [0, 1] \rightarrow \mathbb{R}$ , there exists a real-valued polynomial  $P$  such that for all  $t \in [0, 1]$ ,  $-(10^{-2024} + 2P(t)) \leq f(t) \leq 10^{-2024} - 2P(t)$ .
  - vi. Let  $f \in \mathcal{R}(\alpha)$  in  $[a, b]$  for a given increasing and bounded function  $\alpha$ . Then, there exists a partition  $P$  of  $[a, b]$ ,  $a = x_0 < x_1 < \dots < x_n = b$ , such that for all  $i \in \{1, \dots, n\}$  and for all  $s_i, t_i \in [x_{i-1}, x_i]$  we have that  $\sum_{i=1}^n |f(t_i) - f(s_i)| \Delta \alpha_i \leq 10^{-2024}$ .