
Instructions:

- In case of ambiguity, one has to refer to the ENGLISH version of this exam.
- During the exam you MAY NOT use textbooks, class notes, or any other supporting material apart from the formula sheet given to you.
- Use of calculators is permitted for performing calculations. The only approved calculator are those that do not have graphic or programmable features.
- Start every problem on a new page, and write at the top of the page which problem it belongs to. (But in multiple part problems it is not necessary to start every part on a new page)
- In all of your solutions, give explanations to clearly show your reasoning. Points may be deducted for unclear and wrong argument, even if the final answer is correct.
- Write clearly and legibly.
- Where applicable, indicate your final answer clearly by putting A BOX around it.

Note: There are six problems, some with multiple parts. The problems are not ordered according to difficulty

- (1) (5pt) Compute the degree 3 Taylor polynomial of the function $f(x) = x \ln(1 + (x - 1)^2)$, around the point $x_0 = 1$, and use it to give an approximation of $f(1.1)$.

- (2) Consider the curve in the plane given by the following equation

$$y^2 e^{2x} + C = 3y + x^2,$$

where C is an arbitrary constant.

- (a) (1 pts) Find the value of C such that the curve passes through the point $(0, 3)$.
- (b) (2 pts) With $C = 0$ find the slope of the tangent to the curve in the point $(0, 3)$.
- (c) (1 pt) Find the equation of the tangent to the curve in the point $(0, 3)$.

- (3) Consider the function $f(x) = e^{-x^3+30000x}$.

- (a) (2pts) Find where the function is increasing or decreasing. Find all the critical points and determine their type.
- (b) (1pt) Find the maximum and minimum values of the function on the interval $[-101, -99]$.
- (c) (2 pts) Compute $\lim_{x \rightarrow \pm\infty} f(x)$ and sketch the graph of $f(x)$.

- (4) Compute the following integrals:

(a) (2pt) $\int \left(\frac{\ln(t^5 + 1)}{t^5 + 1} t^4 + 4t^{\frac{7}{5}} \right) dt,$

(b) (3pt) $\int_0^{+\infty} te^{-t} dt.$

(5) Consider the matrix

$$A = \begin{pmatrix} 3 & 0 & 1 \\ 8 & c & 2 \\ c & 6 & 1 \end{pmatrix}$$

- (a) (2 pt) Compute the determinant of A , $|A|$ as a function of c .
- (b) (1 pt) Find all the values of c for which A is not invertible.
- (c) (2 pt) Determine whether the following linear system has 1, 0, or infinitely many solutions. In case there is just one solution, find this.

$$\begin{cases} +2x & +z & = & 1 \\ +8x & +y & +2z & = & 3 \\ x & +6y & +z & = & 3 \end{cases}$$

(6) Consider the two variables function

$$f(x, y) = 48xy - 32x^3 - 24y^2$$

- (a) (2pt) Find all the critical points of $f(x, y)$ and, if possible, determine their type.
- (b) (2pt) Consider now D , the rectangle $0 \leq x \leq 1$ and $0 \leq y \leq 1$. Determine the maximum and minimum values taken by $f(x, y)$ on the *boundary* of the D .
- (c) (1 pt) Determine the minimum and the maximum value of $f(x, y)$ on D .

GOOD LUCK!!!

Svenska Texten

- (1) (5pt) Beräkna det tredje graden Taylors polynom till funktionen $f(x) = x \ln(1+(x-1)^2)$, omkring punkten $x_0 = 1$. Använd det för att approximera $f(1.1)$.

- (2) Antar kurven i planen med ekvation:

$$y^2 e^{2x} + C = 3y + x^2,$$

där C är en godtycklig konstant.

- (a) (1 pts) Hitta värdent till C sådan att kurver går genom punkten med koordinater $(0, 3)$.
- (b) (2 pts) Med $C = 0$ beräkna lutningen till kurven i punkten $(0, 3)$.
- (c) (1 pt) Ange ekvationer of den linjen som är tangenten til kurvan i punkten $(0, 3)$.

- (3) Betrakta funktionen $f(x) = e^{-x^3+30000x}$.

- (a) (2pts) Hitta var funktionen är växande eller avtagende. Hitta all kritiska punkter och bestem deras typ.
- (b) (1pt) Hitta den största och den minsta och minimum värden som funktionen tar i intervallet $[-101, -99]$.
- (c) (2 pts) Beräkna $\lim_{x \rightarrow \pm\infty} f(x)$ och rita grafen till $f(x)$.

- (4) Beräkna de följande integralerna

$$(a) (2pt) \int \left(\frac{\ln(t^5 + 1)}{t^5 + 1} t^4 + 4t^{\frac{7}{5}} \right) dt,$$

$$(b) (3pt) \int_0^{+\infty} te^{-t} dt.$$

- (5) Betrekta matrisen

$$A = \begin{pmatrix} 3 & 0 & 1 \\ 8 & c & 2 \\ c & 6 & 1 \end{pmatrix}$$

- (a) (2 pt) Beräkn determinanten till A , $|A|$ som en funktioner av c .
- (b) (1 pt) Hitta alla värden till c sådana att A int är invertibara.
- (c) (2 pt) Besätm hur många lösningen (1, 0 eller odliga många) har den följande system. Om det finns bara ett, räkna det.

$$\begin{cases} +2x & +z & = & 1 \\ +8x & +y & +2z & = & 3 \\ x & +6y & +z & = & 3 \end{cases}$$

- (6) Betrkata den följnande funktionen i tva variabler

$$f(x, y) = 48xy - 32x^3 - 24y^2$$

- (a) (2pt) Hittal alla den kritiska punkter och, om möjligt, bestäm deras typ.
- (b) (2pt) La D vara rectangeln $0 \leq x \leq 1$ and $0 \leq y \leq 1$. Beräkna den mstörsta och den minsta värden till $f(x, y)$ på gränsen till D .
- (c) (1 pt) Beräkna den största och minsta värden till $f(x, y)$ på D .

Lycka Till!!!