

HOMEWORK SET 4

**HW1.** Let  $X$  be a simply connected space and let  $n \geq 2$ . There is a natural homomorphism

$$\Omega: H^n(X) \rightarrow H^{n-1}(\Omega X)$$

defined by looping  $x: X \rightarrow K(\mathbb{Z}, n)$  and identifying  $\Omega K(\mathbb{Z}, n)$  with  $K(\mathbb{Z}, n-1)$ .

Show that the Serre spectral sequence of the fibration  $\Omega X \rightarrow PX \rightarrow X$  satisfies

- (1)  $E_n^{0, n-1} = \text{im}(\Omega) \subseteq H^{n-1}(\Omega X)$ , and
- (2)  $d_n(\Omega x) \in E_n^{n, 0}$  is equal to the image of  $x$  under the surjection  $H^n(X) \rightarrow E_n^{n, 0}$ .

This shows that the transgression  $\tau = d_n: E_n^{0, n-1} \rightarrow E_n^{n, 0}$  is ‘inverse’ to  $\Omega$  in a certain sense.

**HW2.** Let  $B$  be a simply connected space and let  $n \geq 1$ . Let  $x \in H^{n+1}(B)$  and consider the associated principal fibration

$$K(\mathbb{Z}, n) \rightarrow E \rightarrow B.$$

Show that the  $E_{n+1}$ -page of the Serre spectral sequence with  $\mathbb{Q}$ -coefficients is given by

$$E_{n+1}^{p, q} \cong H^p(B; \mathbb{Q}) \otimes H^q(K(\mathbb{Z}, n); \mathbb{Q}),$$

with differential determined by

$$d_{n+1}(1 \otimes a) = x \otimes 1,$$

where  $a \in H^n(K(\mathbb{Z}, n); \mathbb{Q})$  is the canonical class.

(Hint for both problems: compare with a suitable path space fibration over an Eilenberg–Mac Lane space.)

*Deadline: 2022-11-15. If you have used any resources outside the course literature/lecture notes, please indicate this in your solution. Similarly if you have discussed the problems with another student. Hand in your solutions by e-mail to: alexb@math.su.se*